

When it comes to your hemophilia A treatment

Move beyond the threshold^a

A simple switch to Esperoct[®] can give you high factor levels for longer.^b

^aTrough level goal is 1% for prophylaxis.

^bCompared with standard half-life products.



Discover more at [Esperoct.com](https://www.esperoct.com).

What is Esperoct[®]?

Esperoct[®] [antihemophilic factor (recombinant), glycopegylated-exei] is an injectable medicine to treat and prevent or reduce the number of bleeding episodes in people with hemophilia A. Your healthcare provider may give you Esperoct[®] when you have surgery

- Esperoct[®] is not used to treat von Willebrand Disease

IMPORTANT SAFETY INFORMATION

Who should not use Esperoct[®]?

- You should not use Esperoct[®] if you are allergic to factor VIII or any of the other

ingredients of Esperoct[®] or if you are allergic to hamster proteins

What is the most important information I need to know about Esperoct[®]?

- Do not attempt to do an infusion yourself unless you have been taught how by your healthcare provider or hemophilia treatment center
- Call your healthcare provider right away or get emergency treatment right away if you get any signs of an allergic reaction, such as: hives, chest tightness, wheezing, dizziness, difficulty breathing, and/or swelling of the face

What should I tell my healthcare provider before using Esperoct[®]?

- Before taking Esperoct[®], you should tell your healthcare provider if you have or have had any medical conditions, take any medicines (including non-prescription medicines and dietary supplements), are nursing, pregnant or planning to become pregnant, or have been told that you have inhibitors to factor VIII
- Your body can make antibodies called "inhibitors" against Esperoct[®], which may stop Esperoct[®] from working properly. Call your healthcare provider right

away if your bleeding does not stop after taking Esperoct[®]

What are the possible side effects of Esperoct[®]?

- Common side effects of Esperoct[®] include rash or itching, and swelling, pain, rash or redness at the location of infusion

Please see Brief Summary of Prescribing Information on the following pages.



Novo Nordisk Inc., 800 Scudders Mill Road, Plainsboro, New Jersey 08536 U.S.A.

Esperoct[®] is a registered trademark of Novo Nordisk Health Care AG.

Novo Nordisk is a registered trademark of Novo Nordisk A/S.

© 2020 Novo Nordisk Printed in the U.S.A. US19ESP00185 February 2020

esperoct[®]

*antihemophilic factor (recombinant),
glycopegylated-exei*

esperoct®

antihemophilic factor (recombinant), glycopegylated-exei

Brief Summary information about ESPEROCT® [antihemophilic Factor (recombinant), glycopegylated-exei]

This information is not comprehensive.

- Talk to your healthcare provider or pharmacist
- Visit www.novo-pi.com/esperoct.pdf to obtain FDA-approved product labeling
- Call 1-800-727-6500

Patient Information

ESPEROCT® [antihemophilic factor (recombinant), glycopegylated-exei]

Read the Patient Information and the Instructions For Use that come with ESPEROCT® before you start taking this medicine and each time you get a refill. There may be new information.

This Patient Information does not take the place of talking with your healthcare provider about your medical condition or treatment. If you have questions about ESPEROCT® after reading this information, ask your healthcare provider.

What is the most important information I need to know about ESPEROCT®?

Do not attempt to do an infusion yourself unless you have been taught how by your healthcare provider or hemophilia treatment center.

You must carefully follow your healthcare provider's instructions regarding the dose and schedule for infusing ESPEROCT® so that your treatment will work best for you.

What is ESPEROCT®?

ESPEROCT® is an injectable medicine used to replace clotting Factor VIII that is missing in patients with hemophilia A. Hemophilia A is an inherited bleeding disorder in all age groups that prevents blood from clotting normally.

ESPEROCT® is used to treat and prevent or reduce the number of bleeding episodes in people with hemophilia A.

Your healthcare provider may give you ESPEROCT® when you have surgery.

Who should not use ESPEROCT®?

You should not use ESPEROCT® if you

- are allergic to Factor VIII or any of the other ingredients of ESPEROCT®
- if you are allergic to hamster proteins

If you are not sure, talk to your healthcare provider before using this medicine.

Tell your healthcare provider if you are pregnant or nursing because ESPEROCT® might not be right for you.

What should I tell my healthcare provider before I use ESPEROCT®?

You should tell your healthcare provider if you:

- Have or have had any medical conditions.
- Take any medicines, including non-prescription medicines and dietary supplements.
- Are nursing.
- Are pregnant or planning to become pregnant.
- Have been told that you have inhibitors to Factor VIII.

How should I use ESPEROCT®?

Treatment with ESPEROCT® should be started by a healthcare provider who is experienced in the care of patients with hemophilia A.

ESPEROCT® is given as an infusion into the vein.

You may infuse ESPEROCT® at a hemophilia treatment center, at your healthcare provider's office or in your home. You should be trained on how to do infusions by your hemophilia treatment center or healthcare provider. Many people with hemophilia A learn to infuse the medicine by themselves or with the help of a family member.

Your healthcare provider will tell you how much ESPEROCT® to use based on your weight, the severity of your hemophilia A, and where you are bleeding. Your dose will be calculated in international units, IU.

Call your healthcare provider right away if your bleeding does not stop after taking ESPEROCT®.

If your bleeding is not adequately controlled, it could be due to the development of Factor VIII inhibitors. This should be checked by your healthcare provider. You might need a higher dose of ESPEROCT® or even a different product to control bleeding. Do not increase the total dose of ESPEROCT® to control your bleeding without consulting your healthcare provider.

Use in children

ESPEROCT® can be used in children. Your healthcare provider will decide the dose of ESPEROCT® you will receive.

If you forget to use ESPEROCT®

If you forget a dose, infuse the missed dose when you discover the mistake. Do not infuse a double dose to make up for a forgotten dose. Proceed with the next infusions as scheduled and continue as advised by your healthcare provider.

If you stop using ESPEROCT®

Do not stop using ESPEROCT® without consulting your healthcare provider.

If you have any further questions on the use of this product, ask your healthcare provider.

What if I take too much ESPEROCT®?

Always take ESPEROCT® exactly as your healthcare provider has told you. You should check with your healthcare provider if you are not sure. If you infuse more ESPEROCT® than recommended, tell your healthcare provider as soon as possible.

What are the possible side effects of ESPEROCT®?

Common Side Effects Include:

- rash or itching
- swelling, pain, rash or redness at the location of infusion

Other Possible Side Effects:

You could have an allergic reaction to coagulation Factor VIII products. **Call your healthcare provider right away or get emergency treatment right away if you get any signs of an allergic reaction, such as:** hives, chest tightness, wheezing, dizziness, difficulty breathing, and/or swelling of the face.

Your body can also make antibodies called "inhibitors" against ESPEROCT®, which may stop ESPEROCT® from working properly. Your healthcare provider may need to test your blood for inhibitors from time to time.

These are not all of the possible side effects from ESPEROCT®. Ask your healthcare provider for more information. You are encouraged to report side effects to FDA at 1-800-FDA-1088.

Tell your healthcare provider about any side effect that bothers you or that does not go away.

What are the ESPEROCT® dosage strengths?

ESPEROCT® comes in five different dosage strengths. The actual number of international units (IU) of Factor VIII in the vial will be imprinted on the label and on the box. The five different strengths are as follows:

Cap Color Indicator	Nominal Strength
Red	500 IU per vial
Green	1000 IU per vial
Gray	1500 IU per vial
Yellow	2000 IU per vial
Black	3000 IU per vial

Always check the actual dosage strength printed on the label to make sure you are using the strength prescribed by your healthcare provider.

How should I store ESPEROCT®?

Prior to Reconstitution (mixing the dry powder in the vial with the diluent):

Protect from light. Do not freeze ESPEROCT®.

ESPEROCT® can be stored in refrigeration at 36°F to 46°F (2°C to 8°C) for up to 30 months from the date of manufacture until the expiration date stated on the label.

ESPEROCT® may be stored at room temperature (not to exceed 86°F/30°C), for up to 12 months within the 30-month time period. Record the date when the product was removed from the refrigerator. The total time of storage at room temperature should not exceed 12 months. Do not return the product to the refrigerator.

Do not use this medicine after the expiration date which is on the outer carton and the vial. The expiration date refers to the last day of that month.

After Reconstitution:

The reconstituted (the final product once the powder is mixed with the diluent) ESPEROCT® should appear clear and colorless without visible particles.

The reconstituted ESPEROCT® should be used immediately.

If you cannot use the reconstituted ESPEROCT® immediately, it must be used within 4 hours when stored at or below 86°F (30°C) or within 24 hours when stored in a refrigerator at 36°F to 46°F (2°C to 8°C). Store the reconstituted product in the vial.

Keep this medicine out of the sight and out of reach of children.

What else should I know about ESPEROCT® and hemophilia A?

Medicines are sometimes prescribed for purposes other than those listed here. Do not use ESPEROCT® for a condition for which it is not prescribed. Do not share ESPEROCT® with other people, even if they have the same symptoms that you have.

Revised: 02/2019

ESPEROCT® is a trademark of Novo Nordisk A/S.

For Patent Information, refer to: <http://novonordisk-us.com/patients/products/product-patents.html>

Manufactured by:
Novo Nordisk A/S
Novo Allé
DK-2880 Bagsværd, Denmark

More detailed information is available upon request. Available by prescription only.

For information about ESPEROCT® contact:
Novo Nordisk Inc.
800 Scudders Mill Road
Plainsboro, NJ 08536, USA
1-800-727-6500

© 2019 Novo Nordisk
US19ESP00010 August 2019

