

Move beyond the threshold^a

Esperoct[®] can give you high factor levels for longer.^b



^aOf 1% trough factor levels for standard half-life (SHL) products in adults and adolescents.
^bCompared with SHL products.

WHAT IS ESPEROCT[®]?

Esperoct[®] [antihemophilic factor (recombinant), glycopegylated-exei] is an injectable medicine to treat and prevent or reduce the number of bleeding episodes in people with hemophilia A. Your healthcare provider may give you Esperoct[®] when you have surgery

- Esperoct[®] is not used to treat von Willebrand Disease




Please see Important Safety Information throughout.
Please see accompanying Prescribing Information.

esperoct[®]
*antihemophilic factor (recombinant),
glycopegylated-exei*

Extend half-life beyond the standard

22-hour average half-life in adults^a



Esperoct® 50 IU/kg
Standard rFVIII

rFVIII=recombinant factor VIII.

^aData shown are from 42 adults who received a pharmacokinetic (PK) assessment around the first Esperoct® 50 IU/kg dose.

^bData shown are from a comparison study of 26 previously treated patients (PTPs) 18 years or older who received a 25, 50, or 75 IU/kg dose of their previous SHL product followed by the same dose of Esperoct®. To allow for comparison, all results were adjusted to a 50 IU/kg dose of each product.

Esperoct® is made by taking the existing Novoeight® (rFVIII) molecule and adding PEGylation technology to extend the half-life

IMPORTANT SAFETY INFORMATION

Who should not use Esperoct®?

- You should not use Esperoct® if you are allergic to factor VIII or any of the other ingredients of Esperoct® or if you are allergic to hamster proteins

Please see additional Important Safety Information throughout.

Please see accompanying Prescribing Information.

Level up prophylaxis with a simple dose

High factor levels from one dose to the next

Factor levels stay at or above 3% for

100%
of the time^{c,d}

Factor levels stay at or above 5% for

90%
of the time^{c,e}

Switching made easy with a standard
50 IU/kg dose every 4 days

No dose adjustment needed

**50%
FEWER
INFUSIONS**

if you previously infused every other day

Fewer infusions per year compared with SHL dosing regimens

**40%
FEWER
INFUSIONS**

if you previously infused 3x/week

^cTrough level goal is 1% for prophylaxis.

^dData shown are from a study where 175 previously treated adolescents and adults received routine prophylaxis with Esperoct® 50 IU/kg every 4 days for 76 weeks. Pre-dose factor activity (trough) levels were evaluated at follow-up visits. Mean trough levels for adolescents (12-<18 years) were 2.7 IU/dL.

^eSteady-state factor VIII (FVIII) activity levels were estimated in 143 adults and adolescents using PK modeling.

esperoct®
antihemophilic factor (recombinant), glycopegylated-exei

Stay protected from bleeds

Dose less often^a without sacrificing protection

1.2 Overall bleeds per year^b

0.9 Joint bleeds per year^b


0 Spontaneous bleeds per year^b
Traumatic bleeds per year^b

^aCompared to SHL products, 50% fewer infusions when administered every other day and 40% fewer when administered 3x weekly.

^b175 PTPs with severe hemophilia A received Esperoct[®] 50 IU/kg every 4 days for 76 weeks based on median annualized bleed rates shown.

Prepare for the unexpected

Control for the treatment of bleeding episodes



Dosing for the treatment of bleeding episodes

40 IU/kg for minor to moderate bleeds

50 IU/kg for major bleeds

For moderate to major bleeds, additional dose(s) may be administered every 24 hours

^dData shown are from a study where 12 adult and adolescent PTPs with severe hemophilia chose to be treated on demand and received Esperoct[®] for 532 bleeding episodes.

IMPORTANT SAFETY INFORMATION

What is the most important information I need to know about Esperoct[®]?

- Do not attempt to do an infusion yourself unless you have been taught how by your healthcare provider or hemophilia treatment center
- Call your healthcare provider right away or get emergency treatment right away if you get any signs of an allergic reaction, such as: hives, chest tightness, wheezing, dizziness, difficulty breathing, and/or swelling of the face

Switching made easy

One standard dose makes it easy to switch at any age

65 IU/kg twice weekly

No dose adjustment needed^a

Because FVIII products may be cleared from the body faster in children <12 years, higher and more frequent dosing may be needed.

^aInterval can be adjusted based on individual response to treatment.

IMPORTANT SAFETY INFORMATION


What should I tell my healthcare provider before using Esperoct[®]?

- Before taking Esperoct[®], you should tell your healthcare provider if you have or have had any medical conditions, take any medicines (including non-prescription medicines and dietary supplements), are nursing, pregnant or planning to become pregnant, or have been told that you have inhibitors to factor VIII

Please see additional Important Safety Information throughout.

Please see accompanying Prescribing Information.

Fewer infusions per year compared with SHL dosing regimens



Protection that keeps up with them

Esperoct[®] achieved a

14.3-hour

 average half-life in children^b

85% longer

 half-life compared to SHL^c

Low number of bleeds per year for children aged 0-<12 years^d

2.0

 Overall bleeds per year^d

0

 Joint bleeds
Spontaneous bleeds
Traumatic bleeds^d

^bComparison to prior FVIII product was performed at the beginning of the study in previously treated children. The geometric mean terminal half-life in 23 children aged 0-<12 years was 14.3 hours. Esperoct[®] geometric mean terminal half-life was 14.7 hours in 12 children ages 0-5 and 13.8 hours in 10 children ages 6-11.

^cComparison to prior FVIII product was performed at the beginning of the study in previously treated children. Esperoct[®] half-life was 14.7 hours in 12 children ages 0-6 and 13.8 hours in 10 children ages 6-11.

^dData shown are from a study of 68 previously treated children (34 aged 0-5 and 34 aged 6-11) who received an average dose of approximately 65 IU/kg twice weekly for 26 weeks. Median annualized bleeding rates are shown.



esperoct[®]
antihemophilic factor (recombinant),
glycopegylated-exei

Count on a proven safety profile

270 over **80,000**
previously treated patients (PTPs) infusion days

Safety proven across 5 studies, the largest and longest EHL clinical trial program

- 0 blood clots
- No PEG-related safety concerns
- One PTP with a high-risk gene mutation developed an inhibitor to FVIII^a
 - Similar to the reported rate in patients with severe hemophilia A

EHL=extended half-life; PEG=polyethylene glycol.

^aAn 18-year-old African-American male developed an inhibitor after 93 infusion days of Esperoct[®]. The inhibitor rose to 13.5 Bethesda units and the patient stopped participation in the study. There was no change in efficacy, and the inhibitor eventually went away on its own (without use of immune tolerance induction therapy).

IMPORTANT SAFETY INFORMATION

What should I tell my healthcare provider before using Esperoct[®]? (cont'd)

- Your body can make antibodies called “inhibitors” against Esperoct[®], which may stop Esperoct[®] from working properly. **Call your healthcare provider right away if your bleeding does not stop after taking Esperoct[®]**

Designed to fit into your life

Compact packaging for easy storage



Small kit holds up to a week's worth of factor

Based on a 70 kg person.



Please see additional Important Safety Information throughout.
Please see accompanying Prescribing Information.

esperoct[®]
*antihemophilic factor (recombinant),
glycopegylated-exei*

Flexible on the go

The EHL product with the highest storage temperature for the longest time

Storage Temperature:	Can be stored up to 104°F for up to 3 months	or	Can be stored up to 86°F for up to 12 months
Storage after reconstitution:	Within 4 hours at up to 86°F		

IMPORTANT SAFETY INFORMATION

What are the possible side effects of Esperoct®?

- Common side effects of Esperoct® include rash or itching, and swelling, pain, rash or redness at the location of infusion

Ready in 3 simple steps

A prefilled syringe provides convenient administration in 2 minutes

Attach

Prefilled diluent syringe contains 4 mL of diluent—works with any dose strength


Twist

Adapter connects the syringe and vial with a 25 µm inline particle filter

Mix

After mixing, the reconstituted solution can be administered

5 vial sizes for personalized treatment



Please see additional Important Safety Information throughout.
Please see accompanying Prescribing Information.

esperoct®
antihemophilic factor (recombinant),
glycopegylated-exei

Considerations when switching to Esperoct®

You may be eligible for:

Trial Program

Talk to a NovoSecure™ specialist to find out if you're eligible^a

Product Assistance Program

Apply for the Product Assistance Program by calling 1-844-NOVOSEC (1-844-668-6732) for more information^b

Co-pay Assistance Program

Get help with co-pay costs for Esperoct®^c

^aPatients who have been prescribed a Novo Nordisk hemophilia and rare bleeding disorder product for an FDA-approved indication, and who have commercial insurance, may be eligible to receive a limited supply of free product. Patients who participate in any government, state, or federally funded medical or prescription benefit programs, including Medicare, Medicaid, Medigap, VA, DOD, and TRICARE, including patients who participate in a managed Medicaid program or have Medicaid as secondary insurance, are not eligible to receive product support. Product is provided at no cost to the patient or the HCP, is not contingent on any product purchase, and the patient and HCP must not: (1) bill any third party for the free product, or (2) resell the free product.

^bThe Novo Nordisk Patient Assistance Program (PAP) is administered by NovoSecure™. To qualify for the PAP, patients must demonstrate financial need and must have attempted to find alternative reimbursement. Several factors are considered in evaluating financial need, including cost of living, size of household, and burden of total medical expenses. If the applicant qualifies under the PAP guidelines, a limited supply of the requested medication(s) will be shipped to the patient. Patients who qualify for PAP may be eligible to receive the prescribed Novo Nordisk product, for up to 1 year from the approval date. Product limits vary.

^cNovo Nordisk Hemophilia and Rare Bleeding Disorders Copay/Coinsurance Terms and Conditions: Enrolled patients are eligible for up to \$12,000 in co-pay/coinsurance assistance per calendar year for each NNI hemophilia or rare bleeding disorder product. Assistance is retroactive to 60 days. Patients must be commercially insured and may not be participating in any government, state, or federally funded medical or prescription benefit programs, including Medicare, Medicaid, Medigap, VA, DOD, and TRICARE, including patients who participate in a managed Medicaid program or have Medicaid as secondary insurance. Uninsured, cash-paying patients are not eligible to participate. Patients are eligible to receive co-pay/coinsurance assistance on an annual basis (12 months). Offer good only in the USA, Puerto Rico, Guam, Saipan, and Virgin Islands with participating pharmacies and cannot be redeemed at government-subsidized clinics. Void where taxed, restricted, or prohibited by law. Absent of a change in Massachusetts law, effective July 1, 2019, the Savings Card will no longer be valid for residents of Massachusetts. Patient is responsible for complying with any insurance carrier co-payment disclosure requirements, including disclosing any savings received from this program. Re-confirmation of information may be requested periodically to ensure accuracy of data and compliance with terms. This is not an insurance program. Novo Nordisk reserves the right to rescind, revoke, or amend this offer without notice at any time. Non-medication expenses, such as ancillary supplies or administration-related costs, are not eligible. Must have a current prescription for an FDA-approved indication.

IMPORTANT SAFETY INFORMATION

Who should not use Esperoct®?

- You should not use Esperoct® if you are allergic to factor VIII or any of the other ingredients of Esperoct® or if you are allergic to hamster proteins



Please see additional Important Safety Information throughout.
Please see accompanying Prescribing Information.

esperoct®
*antihemophilic factor (recombinant),
glycopegylated-exei*

Move beyond the threshold^a

Fewer infusions per year compared with SHL for adults and adolescents

- 50% fewer infusions if previously infused every other day
- 40% fewer infusions if previously infused 3x/week

High factor levels in adults and adolescents^b

- At or above 3% for 100% of the time^c
- At or above 5% for 90% of the time^d

Flexible on the go^a

- The only EHL product with stability up to 104°F^e

The largest and longest EHL clinical trial program

^aOf 1% trough factor levels for SHL products in adults and adolescents.

^bData shown are from 42 adults who received a PK assessment around the first Esperoct[®] 50 IU/kg dose.

^cData shown are from a study where 175 previously treated adolescents and adults received routine prophylaxis with Esperoct[®] 50 IU/kg every 4 days for 76 weeks. Pre-dose factor activity (trough) levels were evaluated at follow-up visits. Mean trough levels for adolescents (12-<18 years) were 2.7 IU/dL.

^dSteady-state FVIII activity levels were estimated in 143 adults and adolescents using PK modeling.

^eFor up to 3 months.

IMPORTANT SAFETY INFORMATION

What is the most important information I need to know about Esperoct[®]?

- **Do not attempt to do an infusion yourself unless you have been taught how by your healthcare provider or hemophilia treatment center**

Please see additional Important Safety Information throughout.

Please see accompanying Prescribing Information.

Novo Nordisk Inc., 800 Scudders Mill Road, Plainsboro, New Jersey 08536 U.S.A.

Esperoct[®] is a registered trademark and NovoSecure[™] is a trademark of Novo Nordisk Health Care AG. Novo Nordisk is a registered trademark of Novo Nordisk A/S.

© 2020 Novo Nordisk Printed in the U.S.A. US19ESP00001 January 2020



esperoct[®]
*antihemophilic factor (recombinant),
glycopegylated-exei*